

# Inverse Association of Omentin-1 and Vitamin D with FGF-23 Levels in Prostate Cancer: A Predictive Biomarker Panel

**Khadija A. Sahan\***

Biomedical Informatics College/ University of Information Technology and Communications, Baghdad, Iraq.

## Abstract

**Introduction:** Omentin-1 is typically expressed in vascular endothelial cells, mesothelial cells, airway goblet cells, and adipose tissue stromal cells. Fibroblast growth factor-23 (FGF-23) is primarily secreted by osteocytes, modulates renal phosphate reabsorption and suppresses vitamin D activation. Vitamin D, a fat-soluble vitamin, modulates immune responses, bone metabolism and calcium homeostasis. Prostate cancer remains a major worldwide health concern among males with emerging evidence linking these biomarkers to cancer progression due to their anti-inflammatory, antioxidant and potential anti tumorigenic properties. Objectives: This study aims to determine whether serum levels of omentin-1, vitamin D, and FGF-23, taken together can predict prostate cancer status. **Materials and Methods:** A case-control approach was conducted at a single center Al-Amal Centre/ Medical City of Imam Hussain in Karbala/ Iraq, during December 2022 until June 2023. Serum samples were collected from 30 prostate cancer patients' post-treatment and 30 age-matched (45–70 years) healthy male controls. Serum concentrations of FGF-23, omentin-1, irisin and vitamin D were measured using ELISA. **Results:** The patient group exhibited significantly elevated FGF-23 levels ( $240.8 \pm 55.2$  ng/ml) compared to controls ( $163.1 \pm 22.4$  ng/ml  $p < 0.001$ ). Conversely, the level of irisin; omentin-1; vitamin D was markedly reduced in patients ( $21.9 \pm 3.2$ ,  $10.3 \pm 2.9$ , and  $6.4 \pm 0.7$  ng/ml respectively;  $p < 0.001$ ) relative to controls ( $60.8 \pm 3.5$ ,  $38.8 \pm 6.6$ , and  $19.1 \pm 2.4$  ng/ml respectively). The ROC curve analysis identified serum FGF-23 and omentin-1 and vitamin D as highly accurate molecular indicator for detecting PCa progression (with AUC value; 0.956, 0.958 and 0.938 respectively). **Conclusions:** These circulating factors may be considered as promising diagnostic tools and vitamin D deficiency emerging as a particularly significant risk factor within this study cohort.

**Keywords:** Fibroblast growth factors-23- Irisin- Omentin-1- Prostate cancer- Vitamin D

*Asian Pac J Cancer Care*, **11 (3)**, 441-446

Submission Date: 01/07/2026    Acceptance Date: 03/05/2026

## Introduction

Prostate Cancer (PCa) is the second most frequently tumor diagnosed in male and a top cause of male cancer mortality, representing a critical area of research focus in oncology [1, 2]. The progression of prostate cancer is usually characterized by slow gradual advancement. A silent tumor in aging males, usually remains asymptomatic and undiagnosed during their lifetime [3]. The epidemiology of PCa demonstrates by its association with age. Typically presenting around the seventh decade of life. With nearly two-thirds of deaths occurring in men over 75 [4].

A critical challenge in PCa management is the asymptomatic tumor progression in early and curable

stages, which could delay diagnosis until the tumor progressed, despite diagnosis and therapeutic techniques advancements [5]. In the United States PCa represents a health condition with a population about 3.3 million men with a diagnosis of this tumor, and the mortality rate is approximately 1 in 44 among affected individuals. Conventional PCa diagnosis may lack specificity or lead to overdiagnosis of indolent disease, overtreatment and un-necessary invasive biopsy. Novel biomarkers are essential for more accurate prognosis and the precise identification of aggressive tumors [6].

## Corresponding Author:

Dr. Khadija A. Sahan

Biomedical Informatics College/ University of Information Technology and Communications, Baghdad, Iraq.

Email: dr.Khadija.abbas@uoitc.edu.iq

### *Omentin Gene*

Two isoforms; omentin-1 and omentin-2 are encoded by this gene with 83% amino acids sequence identity. These two isoforms are co-localized on chromosome 1q22-q23. Previous studies have established omentin-1's roles in metabolic regulation and immune responses. Its mature circulating form is a ~120 kDa glycosylated trimer and the bioactive surface is represented by a fibrinogen-like domain (38-82 residues) and a lectin-like domain (37-313 residues) [7].

### *Omentin-1 (Intelectin-1)*

A Multifunctional Adipocytokine with Diagnostic and Therapeutic Potential; Omentin-1, also designated intelectin-1 (ITLN-1), is a 313-amino acid adipocytokine that has emerged as a potential biomarker for various pathological conditions including preeclampsia, inflammatory disorders, polycystic ovarian syndrome, obstructive sleep apnea, and malignancies [8]. This protein is recognized as a diagnostic marker and also as a key acute-phase reactant and represent a promising therapeutic agent [9, 10]. Local ITLN1 expression within the tumor microenvironment may represent a more reliable biomarker of malignant activity than circulating levels. [11]. The positive correlation between omentin-1 and the estradiol-to-testosterone ratio suggests that the reduction in omentin-1 is mechanistically linked to the specific hormonal environment created by ADT [12]. The therapeutic benefit of hormonal ablation in controlling cancers is usually accompanied by clinical issues includes changes in body composition, density of bone mineral, metabolic homeostasis and also sexual functions [13].

### *Fibroblast Growth Factor-23*

FGF-23 is a 251-amino acid endocrine hormone (molecular weight: 32 kDa) encoded by the FGF-23 gene and belongs to the FGF-19/21/23 subfamily. It exhibits mitogenic and anti-apoptotic effects and regulates phosphate homeostasis through modulation of renal phosphate transport [14, 15].

### *FGF-23 in Prostate Cancer*

Pathogenicity and oncogenesis; FGF-23 serves as a potent mitogen in prostate oncogenesis. An autocrine mechanism is observed includes tumor proliferation in prostate, breast and renal carcinoma [16-18]. Its primary site of synthesis is the osteocyte and play a principal physiological role in modulating vitamin D and phosphate levels [19]. The effect of this protein includes two complementary pathways; first, the direct effect on proximal renal tubules by suppressing of phosphate reabsorption [20]. Second, through indirect inhibition of renal 1  $\alpha$ -hydroxylase activity which diminishes the production of calcitriol [21]. The dual effect creates a hypo-phosphatemic condition. This was supported by preclinical studies, including human tumor specimen, in vitro models and animal systems demonstrate that vitamin D deficiency with FGF-23 overexpression induce prostate carcinogenesis [22].

### *The role of Irisin*

Irisin is a glycoprotein composed of 112 amino acids secreted from skeletal muscle and fat tissue [23]. It functions in browning of white adipose tissues. enhancing their metabolic activity [24]. Clinical oncology evidences represent an important irisin expression pattern, dysregulation of irisin expression across several major cancers including breast cancer, obesity related malignancies, hormone dependent cancers and prostate carcinoma [25,26].

### *Vitamin D; Metabolism and clinical importance*

Vitamin D is found in 2 primary forms. The first form is ergocalciferol (D2). The second form is cholecalciferol (D3) [27]. The activation of both forms requires 25-hydroxylation in the liver [28], but cholecalciferol (D3) demonstrate some advantages in bioavailability and efficacy. So, the serum concentration of cholecalciferol (D3) considered as the main biomarker for vitamin D deficiency indication [29]. Vitamin D deficiency is epidemiologically associated with increased cancer risk and adverse oncological outcomes, with marked racial disparities compounding its role as a significant public health challenge [30-32].

"The hypothesis of this study is that a combined panel of omentin-1, FGF-23, and vitamin D can more accurately predict prostate cancer status than individual biomarkers."

### *Study Design and Population*

This study employed a case-control study, in which serum samples from both PCa patients and healthy controls were obtained and analyzed at Al Amal Oncology Centre in Karbala/ Iraq, within December 2022 and June 2023. Serum concentrations of irisin, omentin-1; FGF-23 and vitamin D have been quantified by the use of ELISA. Analyses were employed by the use of commercial ELISA kits from Cloud-Clone Corp. (USA) were employed: SEN576Hu (for irisin), SEA933Hu (for omentin-1), SEA746Hu (for FGF-23), and CEA920Ge (for vitamin D) according to the manufacturer's instructions.

Inclusion criteria: The study comprised a total of 60 participants, including 30 prostate cancer patients receiving hormonal therapy with Zoladex and thirty apparently healthy controls. Both cases and controls were selected within the age range of 45 to 70 years.

Irisin was investigated as an exploratory, mechanistic biomarker within the panel.

Exclusion criteria: for prostate cancer patients and controls included a history of diabetes, thyroid disorders, or cardiovascular diseases to minimize potential confounding effects.

The Ministry of Health's Kerbala Directorate of Health's study committee granted ethical permission (Ref: 206 in 27/November 2022).

### *Statistics and data analysis*

Using T-test to compare between studied groups. With p-value of < 0.05; significant statistical association. Pearson's correlation values were obtained by use of a simple linear regression model to assess associations

between variables. All statistical analyses and data management were conducted using ROC curve and SPSS software, version 28.0 (Released 2021; IBM Corp. USA).

For all inferential tests a two-tailed approach was used [33,34].

A priori power analysis indicated that a sample size of N=30 per group provides 80% power ( $\alpha=0.05$ ) to detect a medium-to-large effect size (Cohen's  $d \geq 0.7$ ). To account for multiple comparisons, a Bonferroni correction was considered; however, all reported p-values ( $< 0.001$ ) remained statistically significant after adjustment.

## Results

As shown in Table 1, the mean serum irisin level was significantly lower within patient group ( $21.9 \pm 3.2$ ) ng/ml in comparison to the healthy control category ( $60.8 \pm 3.5$ ) ng/ml,  $p < 0.001$ ). Similarly, the average level in serum of FGF-23 (ng/ml) in prostate cancer group undergoing hormone therapy was significantly elevated ( $240.8 \pm 55.2$ ) compared to apparently healthy controls ( $163.1 \pm 22.4$ ) with ( $p < 0.001$ ). The mean serum concentration for omentin-1 (ng/ml) were also markedly reduced in PCa patients ( $10.3 \pm 2.9$ ) relative to the apparently healthy control group ( $38.8 \pm 6.6$ ) with ( $p < 0.001$ ). Furthermore, the average level for vitamin D level (ng/ml) was significantly lower in patients ( $6.4 \pm 0.7$ ) than in controls ( $19.1 \pm 2.4$ ), ( $p < 0.001$ ). Body mass index in patients is ranging from overweight to mild obesity (BMI:  $26.0 - 29.0$  kg/m<sup>2</sup>; Overweight - Obese class I) and for healthy control group is ranging from normal to overweight (BMI:  $24.5 - 27.0$  kg/m<sup>2</sup>; Normal - Overweight). Tables 2 and 3 illustrate the relationship between FGF-23 and omentin-1 with other measured parameters in PCa patients receiving hormonal therapy. Table 2 represents a non-significant negative relationship between FGF-23 and irisin, and a non-significant positive association with vitamin D. However, a statistically significant positive

correlation was observed between FGF-23 and omentin-1. The data in Table 3 indicate a statistically significant positive correlation between omentin-1 and vitamin D values. The correlation between omentin-1 and irisin was positive but not significant.

**ROC Curve Analysis:** This curve was employed to find out the discriminatory effect of irisin; omentin-1; FGF-23 and vitamin D as important biomarkers (Table 4). This curve graphically explained the inverse relationship between sensitivity and specificity for these biomarkers. The Area Under the Curve (AUC) values for ROC analysis was:

Irisin: moderate discriminative ability (AUC = 0.733; 95% CI: 0.644–0.824). FGF-23: excellent diagnostic accuracy (AUC = 0.956; 95% CI: 0.993–0.988), omentin-1: (AUC = 0.958; 95% CI: 0.929–0.990) and vitamin D: (AUC = 0.938; 95% CI: 0.902–0.963).

These values indicate that irisin has a moderate discriminative power. Omentin-1, FGF-23 and vitamin D possess excellent discriminative power between control and patients and predicting disease progression and treatment response. For FGF-23, a cutoff of 252.65 ng/ml, sensitivity (95.3%), specificity (99.8%), suggesting it as a suitable biomarker for diagnostic conformation, although a small part of prostate cancer cases ( $\approx 4.7\%$ ) may be missed.

Optimal cutoff values were determined using the Youden index to maximize the combined sensitivity and specificity.

## Discussion

The decreased level of irisin in studied PCa patients is consistent with Tekin et al. [35], they identified irisin as an important key regulator for PCa cell proliferation. Another study reported a significant low level of irisin in PCa patients, suggesting irisin as a promising biomarker which could enhance the diagnosis and prognosis alongside prostate-specific antigen (PSA) [36]. In treated patients,

Table 1. Circulating Levels of Irisin, Omentin-1, FGF-23 and Vit. D (ng/ml) in Control and PCa Groups

Parameters (ng/ml)	Mean $\pm$ SD		P-value
	Controls (NO.30)	Patients (No.30)	
Irisin	60.8 $\pm$ 3.5	21.9 $\pm$ 3.2	0.0001
Omentin-1	38.8 $\pm$ 6.6	10.3 $\pm$ 2.9	0.0001
FGF-23	163.1 $\pm$ 22.4	240.8 $\pm$ 55.2	0.0001
Vit. D	19.1 $\pm$ 2.4	6.4 $\pm$ 0.7	0.0001

Table 2. Statistical Relationship between FGF-23 with other 3 Parameters

FGF-23	Factors	Irisin	Omentin-1	Vit. D
	R (Pearson)		-0.011	0.473
	P	0.954	0.008	0.393

Table 3. Statistical Relationship between Omentin-1 with (Irisin and Vit. D)

Omentin-1	Factor	Irisin	Vit. D
	R (person)		0.055
	P	0.774	0.0001

Table 4. ROC Curve and AUC Analysis for Irisin, Omentin-1, FGF-23 and Vit. D.

Variables	Area under the curve	Sensitivity	Specificity	95% C.I		Cut off value
				L.B.	U.B.	
Irisin	0.733	0.852	0.718	0.644	0.824	21.794
FGF-23	0.956	0.953	0.998	0.993	0.988	252.65
Omentin-1	0.958	0.935	0.962	0.929	0.99	25.787
Vit. D	0.938	0.927	0.952	0.902	0.963	15.51

the decline in irisin circulatory levels reflects a catabolic consequence induced by therapy, characterized by weight loss and decline in primary tissue source (muscle mass loss) [37]. Consistent with current data, Teishima et al. confirmed that FGF-23 levels are increased in PCa patients [38]. The elevated FGF-23 serum levels in PCa patients in this study is consistent with Hussein et al. which noted that androgen-deprivation therapy in PCa patients can elevate FGF-23, through the induction of osteoporosis [39]. The suppression of vitamin D synthesis by FGF2-3 regulated by downregulation of renal enzyme 1 $\alpha$ -hydroxylase (CYP27B1), create a physiological environment for inverse relationship between increased FGF-23 and decreased 1,25-dihydroxyvitamin D [1,25 (OH)<sub>2</sub>D] observed in PCa patients serum levels [40]. Beyond the classical function of vitamin D and its role in mineral homeostasis, it also recognized as a key regulator in carcinogenesis. Recent evidences from epigenomic, transcriptomic and proteomic studies have proved its involvement in cancer cell regulation, self renewal and apoptosis. Beside direct role and cellular effects, vitamin D display important immunomodulatory capacities in tumor microenvironment and anti-neoplastic effects. Low circulating levels of this vitamin are correlated with increased cancer risk. Therefore, vitamin D supplementation, as monotherapy or conjugated with immunotherapy represent a promising strategy for oncological outcome improvement [41,42]. Our findings demonstrate that a combined panel of vitamin D, omentin-1, and FGF-23 offers promising diagnostic potential. Vitamin D deficiency, supported by evidence of impaired activation by elevated FGF-23, is a central feature of our patient cohort. This mechanism is consistent with Layne., et al and suggests that vitamin D supplementation warrants investigation as a therapeutic adjunct [43].

In conclusion, this study suggests that a combined panel of vitamin D, omentin-1, and FGF-23 may serve as a diagnostic tool to complement PSA in prostate cancer detection. However, prospective validation in independent cohorts and assessment of clinical utility in risk stratification are required before clinical implementation.

#### Declarations

The authors declare no conflicts of interest.

#### Funding

This research received no external funding.

#### Clinical trial registration

Not applicable.

#### Conflicts of interest/Competing interests

Authors declare that they have no conflicts of interest.

#### Availability of data and material

The data sets used and/or analyzed during the current study are available from the corresponding authors per reasonable request.

#### Code availability

No custom code was used in this study.

#### Authors' contributions

The author contributed to the conception and design of the study, data collection, data analysis, and drafting and final revision of the manuscript. The author approved the final version for submission.

#### Ethics approval

This study was approved by the Ethic Committee of the Ministry of Health / Karbala Health Directorate; No. 206 Date: 27/11/2022.

#### Consent to participate

Written informed consent was obtained from all participants prior to their inclusion in the study, and the study was conducted in accordance with the principles of the Declaration of Helsinki.

#### Consent for publication

Written informed consent was obtained from all participants, and the trial was conducted in accordance with the Declaration of Helsinki.

#### Originality Declaration for Figures and Tables

All tables included in this manuscript are original and have been created by the authors specifically for the purposes of this study. No previously published or copyrighted images have been used. The authors confirm that all graphical elements, illustrations, and visual materials were generated from the data obtained in the course of this research or designed uniquely for this manuscript.

## Acknowledgments

We extend our heartfelt gratitude to all the patients who generously contributed to this study. We are particularly thankful to the dedicated nursing staff of the Al-Amal Centre/ Medical City of Imam Hussain in Karbala/ Iraq, whose invaluable assistance and compassionate care greatly supported the research process.

*Declaration on generative AI and AI-assisted technologies in the writing process*

AI paraphrasing tools were used.

## References

- Varaprasad GL, Gupta VK, Prasad K, Kim E, Tej MB, Mohanty P, Verma HK, et al. Recent advances and future perspectives in the therapeutics of prostate cancer. *Experimental Hematology & Oncology*. 2023 09 22;12(1):80. <https://doi.org/10.1186/s40164-023-00444-9>
- Al-qazzaz A, Altaie AF. The Role of Omentin-1 and Fibroblast Growth Factor-23 in Iraqi Patients with Prostate Cancer during Chemotherapy. *Journal of the Faculty of Medicine Baghdad*. 2024 Oct 01;66(3):254-259. <https://doi.org/10.32007/jfacmedbaghdad.6632256>
- Bleyer A, Spreafico F, Barr R. Prostate cancer in young men: An emerging young adult and older adolescent challenge. *Cancer*. 2020 01;126(1):46-57. <https://doi.org/10.1002/cncr.32498>
- Badal S, Aiken W, Morrison B, Valentine H, Bryan S, Gachii A, Ragin C. Disparities in prostate cancer incidence and mortality rates: Solvable or not?. *The Prostate*. 2020 01;80(1):3-16. <https://doi.org/10.1002/pros.23923>
- James LJ, Wong G, Craig JC, Hanson CS, Ju A, Howard K, et al. Men's perspectives of prostate cancer screening: A systematic review of qualitative studies. *PloS one*. 2017 Nov 28;12(11). <https://doi.org/10.1371/journal.pone.0188258> PMID 29182649
- National Cancer Institute. SEER Cancer Stat Facts: Prostate Cancer. Accessed at <https://seer.cancer.gov/statfacts/html/prost.html> on June 23, 2023. <https://seer.cancer.gov/statfacts/html/prost.html>.
- Ahmed HS, Salman ED. Role of Serum Omentin-1 and Bone Metabolism Markers in Osteoporosis among Postmenopausal Women. *Annals of the Romanian Society for Cell Biology*. 2021;25(6):8317-27. <https://annalsofscb.ro/index.php/journal/article/view/7025>.
- Tao M, Yan W, Chen C, Tang M, Zhao X, Feng Q, Fei X, Fu Y. Omentin-1 ameliorates experimental inflammatory bowel disease via Nrf2 activation and redox regulation. *Life Sciences*. 2023 09;328:121847. <https://doi.org/10.1016/j.lfs.2023.121847>
- Kadium TE, Alrubaie A, Ghanim SAM. The Link between Serum Omentin Level and Insulin Resistance Biomarkers, Lipid Profile, and Atherogenic Indices in Iraqi Obese Patients. *Baghdad Science Journal*. 2023 02 01;20(1):0074. <https://doi.org/10.21123/bsj.2022.6535>
- Alkhader RAY, Ali KAW, Al-Kharasani AMR. Irisin Levels Related to Glycosylated Hemoglobin in Type 2 Diabetic Obese Women. *Iraqi Journal of Science*. 2020 09 29;2150-2155. <https://doi.org/10.24996/ij.s.2020.61.9.1>
- Sekhoacha M, Riet K, Motloung P, Gumenuku L, Adegoke A, Mashele S. Prostate Cancer Review: Genetics, Diagnosis, Treatment Options, and Alternative Approaches. *Molecules*. 2022 09 05;27(17):5730. <https://doi.org/10.3390/molecules27175730>
- A B, L S. Serum Omentin Levels in Patients with Prostate Cancer and Associations with Sex Steroids and Metabolic Syndrome. *Journal of clinical medicine*. 2020 04 20;9(4). <https://doi.org/10.3390/jcm9041179> PMID 32326011
- Reckell T, Nguyen K, Phan T, Crook S, Kostelich EJ, Kuang Y. Modeling the synergistic properties of drugs in hormonal treatment for prostate cancer. *Journal of Theoretical Biology*. 2021 04;514:110570. <https://doi.org/10.1016/j.jtbi.2020.110570>
- Ali IA, Ali SH. Impact of Osteocalcin Level on Vascular Calcification in Type 2 Diabetics in Relation to Fibroblast Growth Factor-23(FGF-23). *Iraqi Journal of Pharmaceutical Sciences (P-ISSN: 1683 - 3597 , E-ISSN : 2521 - 3512)*. 2018 Dec 06;42-54. <https://doi.org/10.31351/vol27iss2pp42-54>
- Thajeel TS, Rasheed MK, Sudany A, Hanon MJB. Evaluation of cystatin c and fibroblast growth factor 23 in type 2 diabetic patients and its association with early nephropathy. *Biochemical & Cellular Archives*. 2021;21(1). <https://connectjournals.com/03896.2021.21.1429>.
- Ameen ZA, Ali SH. Effects of Aldosterone, Osteoprotegerin and Fibroblast Growth Factor-23 and Some Biochemical Markers in Chronic Kidney Disease Patients (Stage II-IV) among Patients with or without Cardiovascular Events. *Iraqi Journal of Pharmaceutical Sciences (P-ISSN: 1683 - 3597 , E-ISSN : 2521 - 3512)*. 2018 Dec 07;150-158. <https://doi.org/10.31351/vol27iss2pp150-158>
- Colangelo L, Sonato C, Cipriani C, Pepe J, Farinacci G, Palmisano B, Occhiuto M, et al. Occipital bone and tumor-induced osteomalacia: a rare tumor site for an uncommon paraneoplastic syndrome. *Archives of Osteoporosis*. 2023 07 12;18(1):94. <https://doi.org/10.1007/s11657-023-01305-y>
- Al-Khaykanee() AM, Abdel-Rahman AA, Essa A, Gadallah AA, Ali BH, Al-Aqar AA, Badr EA, Shehab-Eldeen S. Genetic polymorphism of fibroblast growth factor receptor 2 and trinucleotide repeat-containing 9 influence the susceptibility to HCV-induced hepatocellular carcinoma. *Clinics and Research in Hepatology and Gastroenterology*. 2021 Nov;45(6):101636. <https://doi.org/10.1016/j.clinre.2021.101636>
- Kashash SA, Hameed AS. Evaluating the Fibroblast Growth Factor-23 and Phosphate in Iraqi Patients with Acromegaly. *Baghdad Science Journal*. 2024 03 04;21(3):1037. <https://doi.org/10.21123/bsj.2023.8814>
- Kuloğlu T, Artaş G, Yardim M, Sahin I, Aydin Y, Beyoğlu N, Özeran İH, et al. Immunostaining characteristics of irisin in benign and malignant renal cancers. *Biotechnic & Histochemistry*. 2019 08 18;94(6):435-441. <https://doi.org/10.1080/10520295.2019.1586998>
- Nakatani S, Nakatani A, Mori K, Emoto M, Inaba M, Razaque MS. Fibroblast Growth Factor 23 as Regulator of Vitamin D Metabolism. *Phosphate Metabolism*. 2022;1362:47-54. [https://doi.org/10.1007/978-3-030-91623-7\\_6](https://doi.org/10.1007/978-3-030-91623-7_6)
- Mizuno Y, Ishida T, Kugimiya F, Takai S, Nakayama Y, Yonemitsu K, Harada E. Deterioration of Phosphate Homeostasis Is a Trigger for Cardiac Afterload — Clinical Importance of Fibroblast Growth Factor 23 for Accelerated Aging —. *Circulation Reports*. 2023 01 10;5(1):4-12. <https://doi.org/10.1253/circrep.CR-22-0124>
- Abdul Sattar A, T. AL-Sa'adi W. Evaluation of Adiponectin, Irisin, Vitamin D Levels and Their Relation in Iraqi Patients with Non alcoholic Fatty Liver Disease. *Ibn AL-Haitham Journal For Pure and Applied Sciences*. 2023 04 20;36(2):241-250. <https://doi.org/10.30526/36.2.3043>
- Al-Aridhi DTN, Allehibi KIH, Al-Sharifi ZA, Al Quraishi

- MJM-LU. Serum Levels of Novel Biochemical Marker (Irisin) in Relation to the Duration of Type 2 Diabetes & in Cases of Type 2 diabetes with Coronary Artery Disease in Iraqi Patients Aged (40-60 year). *Medico-Legal Update*. 2020;20(1). <https://doi.org/10.37506/v20%2Fi1%2F2020%2Fmlu%2F194390>
25. Cebulski K, Piotrowska A, Kmiecik A, Haczekiewicz-Leśniak K, Ciesielska U, Grzegorzółka J, Jabłońska K, et al. The Role of Irisin/FNDC5 Expression and Its Serum Level in Breast Cancer. *International Journal of Molecular Sciences*. 2023 05 11;24(10):8628. <https://doi.org/10.3390/ijms24108628>
  26. Alshaqiti KH, Alomar SF, Alzoman N, Almomen A. Irisin Induces Apoptosis in Metastatic Prostate Cancer Cells and Inhibits Tumor Growth In Vivo. *Cancers*. 2023 08 07;15(15):4000. <https://doi.org/10.3390/cancers15154000>
  27. Rondanelli M, Moroni A, Zese M, Gasparri C, Riva A, Petrangolini G, Perna S, Mazzola G. Vitamin D from UV-Irradiated Mushrooms as a Way for Vitamin D Supplementation: A Systematic Review on Classic and Nonclassic Effects in Human and Animal Models. *Antioxidants*. 2023 03 16;12(3):736. <https://doi.org/10.3390/antiox12030736>
  28. Hussein TA, Shafeeq NK, Hussain NA. Comparison of 1, 25-Dihydroxyvitamin D3 and Glutathione Reductase Levels in Thyroid Gland Patients. *Pakistan Journal of Medical and Health Sciences*. 2021;15(1):224 - 9. <https://pjmhsonline.com/2021/jan/224.pdf>.
  29. Salman Jasim H, Khalid Shafeeq N, Abass EAA. Vitamin D Level and its Relation with the Newly Diagnosed Diabetic Neuropathy in Women with Hypothyroidism. *Archives of Razi Institute*. 2022 06 30;77(3):1139-1145. <https://doi.org/10.22092/ari.2022.357389.2029>
  30. Ismail NH, Mussa A, Al-Khreisat MJ, Mohamed Yusoff S, Husin A, Johan MF, Islam MA. The Global Prevalence of Vitamin D Deficiency and Insufficiency in Patients with Multiple Myeloma: A Systematic Review and Meta-Analysis. *Nutrients*. 2023 07 20;15(14):3227. <https://doi.org/10.3390/nu15143227>
  31. Al-Mendalawi MD. Correlation of Vitamin D3 Levels with Disease Severity in Psoriasis Patients – A Case-Control Study. *Indian Journal of Dermatology*. 2023 05;68(3):356-356. [https://doi.org/10.4103/ijd.ijd\\_211\\_23](https://doi.org/10.4103/ijd.ijd_211_23)
  32. Moslem MN, Altaie AF. Correlation study of (IL-17, IL-17RA) and some biochemical parameters with patients of Iraq colorectal cancer after and before taking a dose of chemotherapy. 2020;:020295. <https://doi.org/10.1063/5.0000066>
  33. Mat Roni S, Djajadikerta HG. Data Analysis with SPSS for Survey-based Research. 2021. <https://doi.org/10.1007/978-981-16-0193-4>
  34. Sahan K, Sahan ZA, Qazzaz HA. Predictive Power of Multiplex Serum Biomarkers in Breast Cancer: A Comprehensive ROC Curve Analysis. *Iraqi Journal of Cancer and Medical Genetics*. 2025 Dec 30;18(2). <https://doi.org/10.29409/zvqcs091>
  35. Tekin S, Erden Y, Sandal S, Yilmaz B. Is irisin an anticarcinogenic peptide?. *Medicine Science*. 2015;4(2):2172–80. <https://doi.org/https://doi.org/10.5455/medscience.2014.03.8210>
  36. Aslan R, Alp H, Eryılmaz R, Huyut Z, Sevim M, Araz Ş. Can the irisin be a biomarker for prostate cancer? A case control study. 2020;. <https://doi.org/https://doi.org/10.31557%2FAPJCP.2020.21.2.505>
  37. Huh J, Panagiotou G, Mougios V, Brinkoetter M, Vamvini M, Schneider B. FNDC5 and irisin in humans: I. Predictors of circulating concentrations in serum and plasma and II. mRNA expression and circulating concentrations in response to weight loss and exercise. *Metabolism*. 2012;61(12):1725–38. <https://doi.org/10.1016/j.metabol.2012.09.002>
  38. Teishima J, Hayashi T, Nagamatsu H, Shoji K, Shikuma H, Yamanaka R. Fibroblast growth factor family in the progression of prostate cancer. *Journal of Clinical Medicine*. 2019;8(2). <https://doi.org/10.3390/jcm8020183>
  39. Hussain A, Tripathi A, Pieczonka C, Cope D, McNatty A, Logothetis C. Bone health effects of androgen-deprivation therapy and androgen receptor inhibitors in patients with nonmetastatic castration-resistant prostate cancer. *Prostate cancer and prostatic diseases*. 2021;24(2):290–300. <https://doi.org/10.1038/s41391-020-00296-y>
  40. Carlberg C, Raczky M, Zawrotna N. Vitamin D: A master example of nutrigenomics. *Redox Biology*. 2023 06;62:102695. <https://doi.org/10.1016/j.redox.2023.102695>
  41. Seraphin G, Rieger S, Hewison M, Capobianco E, Lisse TS. The impact of vitamin D on cancer: A mini review. *The Journal of Steroid Biochemistry and Molecular Biology*. 2023 07;231:106308. <https://doi.org/10.1016/j.jsbmb.2023.106308>
  42. Sahan KA, Sahan ZA, Turki TG. A Computational Analysis of Four RETNSNPs in Iraqi Women with Breast Cancer: A Cross-Sectional Study. *Journal of the Faculty of Medicine Baghdad*. 2025 Oct 01;67(3):359-367. <https://doi.org/10.32007/jfacmedbaghdad3148>
  43. Layne TM, Rothstein JH, Song X, Andersen SW, Benn EK, Sieh W, Klein RJ. Vitamin D-related genetic variants and prostate cancer risk in Black men. *Cancer Epidemiology*. 2025 04;95:102742. <https://doi.org/10.1016/j.canep.2025.102742>



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.